

220 mm

140 mm

<p>Scheduling status S1</p> <p>Proprietary name and dosage form</p> <p>Andolex - C</p> <p>ORAL GEL (Gel)</p> <p>PEEL HERE TO REVEAL PRODUCT INFORMATION</p>	
<p>PRINT FREE AREA FOR GLUING</p> <p>PRINT FREE AREA FOR GLUING</p>	
<p>COMPOSITION Each 1 g contains: Benzylamine hydrochloride 10,0 mg Cetylpyridinium chloride 1,0 mg Other ingredients are glycerol, natrosol 250 HHX pharma, peppermint oil crystal 2110, polysorbate 20, purified water and sodium saccharin. SUGAR FREE.</p> <p>PHARMACOLOGICAL CLASSIFICATION A 16.4 Naso-pharyngeal and bucco-pharyngeal antiseptics.</p> <p>PHARMACOLOGICAL ACTION Benzylamine hydrochloride has local analgesic and anti-inflammatory properties. Cetylpyridinium chloride is a quaternary pyridinium antiseptic with bactericidal and fungicidal activity.</p> <p>INDICATIONS Temporarily relieves painful inflamed conditions of the mouth, including mouth and denture ulcers and sore gums.</p> <p>CONTRAINDICATIONS Patients with known hypersensitivity to benzylamine or cetylpyridinium chloride. ANDOLEX-C ORAL GEL is not recommended in children under 6 years of age. Refer to PREGNANCY AND LACTATION</p> <p>WARNINGS AND SPECIAL PRECAUTIONS If a sore mouth is caused or complicated by a bacterial infection, appropriate antibacterial therapy should be considered.</p> <p>PREGNANCY AND LACTATION The safety of benzylamine hydrochloride has not been established in pregnancy and lactation.</p> <p>DOSAGE AND DIRECTIONS FOR USE 1. Apply approximately 1 cm of gel with finger. 2. Gently massage into sore area. 3. Do not eat or drink for 15 minutes. 4. Apply every 2 to 3 hours up to a maximum of 12 times/day. If symptoms persist, see your doctor or dentist.</p> <p>SIDE EFFECTS The most commonly reported reaction is oral numbness. Burning or stinging sensation has been reported. Other local adverse effects include dryness of mouth or thirst, tingling, warm feeling in mouth and altered sense of taste.</p>	<p>Immune System Disorders <i>Frequency unknown:</i> Serious allergic reaction (anaphylactic shock), signs of which may include difficulty breathing, chest pain or chest tightness, and/or feeling dizzy/faint, severe itching of the skin or raised lumps on the skin, swelling of the face, lips, tongue and/or throat, and which may be potentially life-threatening.</p> <p>KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT See SIDE EFFECTS and WARNINGS AND SPECIAL PRECAUTIONS. Adverse central nervous system effects have been reported following overdosage with high doses of benzylamine hydrochloride in solution form. Nausea and vomiting may also occur. There is no specific antidote for benzylamine and treatment should be symptomatic and supportive.</p> <p>IDENTIFICATION. Clear, colourless, peppermint flavoured gel.</p> <p>PRESENTATION White aluminium laminated tubes containing 10 g.</p> <p>STORAGE INSTRUCTIONS Store at or below 30 °C. KEEP OUT OF REACH OF CHILDREN Keep well closed.</p> <p>REGISTRATION NUMBER 33/16.4/0285</p> <p>NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION iNova Pharmaceuticals (Pty) Ltd 15e Riley Road, Bedfordview 2007</p> <p>DATE OF PUBLICATION OF THE PACKAGE INSERT 26 April 2002</p> <p>BOTSWANA Scheduling Status: 3 Pharmacological classification: ATC: A01A D02 Stomatological preparations: other agents for local oral treatment Licence Number: BOT 0200516</p> <p>NAMIBIA Scheduling Status: NS1 Registration Number: 06/16.4/0095</p> <p>inova pharmaceuticals LT015503 LT049.08.17</p>

<p>Skeduleringstatus S1</p> <p>Eiendomsnaam en doseervorm</p> <p>Andolex - C</p> <p>ORAL GEL (Gel)</p>		
<p>SAMESTELLING Elke 1 g bevat: Bensidamienhydrochloried 10,0 mg Setielpiridiniumchloried 1,0 mg Ander onaktiewe bestanddele sluit in gliserol, natrosol 250 HHX pharma, pepermentolie kristalle 2110, polisorbataat 20, gesuiwerde water and natriumsakkarien. SUIKER VRY.</p> <p>FARMAKOLOGIESE KLASSEKASIE A 16.4 Neus-antiseptika en mond-en-keelantiseptika.</p> <p>FARMAKOLOGIESE WERKING Bensidamienhydrochloried het plaaslike analgetiese en anti-inflammatoriese eienskappe. Setielpiridiniumchloried is 'n kwaternêre antiseptikum met kiemdodende en swamdodende aktiwiteit.</p> <p>INDIKASIES Tydelike verligting van pynlike geïntlammeerde kondisies van die mond, insluitende mond en kunstand ulserasies en seer tandvleise.</p> <p>KONTRA-INDIKASIES Pasiënte met bekende hipersensitiwiteit vir bensidamienhydrochloried of setielpiridiniumchloried. ANDOLEX-C ORAL GEL word nie aanbeveel vir kinders jonger as 6 jaar oud nie. Verwys na SWANGESKAP EN LAKTASIE.</p> <p>WAARSKUWINGS EN SPESIALE VOORSORGMATREËLS Indien 'n seer mond veroorsaak is of gekompliseer word deur 'n bakteriële infeksie, moet toepaslike antibakteriële terapie oorweeg word.</p> <p>SWANGESKAP EN LAKTASIE Die veiligheid van bensidamienhydrochloried is nog nie vasgestel in swangerskap en laktasie nie.</p> <p>DOSIS EN GEBRUIKSAANWYSINGS 1. Wend ongeveer 1 cm gel met die vinger aan. 2. Masseer sagkuns in die seer area in. 3. Moet nie eet of drink vir 15 minute nie. 4. Wend elke 2 tot 3 ure aan tot 'n maksimum van 12 keer/dag. Indien simptome volhou, raadpleeg u dokter of tandarts.</p>		<p>NEWE-EFFEKTE Die mees algemene reaksie is 'n dooierige gevoel in die mond. 'n Branderige- of prikkelsensasie is al aangemeld. Ander plaaslike newe-effekte sluit in droogheid van die mond of dors, prikkeling, 'n warm gevoel in die mond en 'n veranderde sin van smaak.</p> <p>Immunsisteem Verstourings <i>Frekwensie onbekend:</i> Ernstige allergiese reaksie (anafylaktiese skok), tekens sluit moeilike asemhaling, borspyn of bors wat toetrek, en / of duiseligheid/flou gevoel, erge jeuk van die vel of ophewe knoppe op die vel, swelling van die gesig, lippe, tong en / of keel in, en kan potensieel lewensgevaarlik wees.</p> <p>BEKENDE SIMPTOME VAN OORDOSERING EN BESONDERHEDE VAN DIE BEHANDELING DAARVAN Sien NEWE-EFFEKTE en WAARSKUWINGS en SPESIALE VOORSORGMATREËLS. Newe-effekte van die sentrale senuweestelsel is vermeld met oordosering van bensidamienhydrochloried in oplossing. Naarheid en braking kan voorkom. Daar is geen spesifieke teenmiddel vir bensidamienhydrochloried nie, en behandeling behoort simptome en ondersteunend te wees.</p> <p>IDENTIFIKASIE Helder, kleurlose, peppermint geurde gel.</p> <p>AANBIEDING Wit aluminium gelamineerde buise wat 10 g bevat.</p> <p>BERGINGSAAANWYSINGS Bewaar teen of benede 30 °C. HOU BUITE BEREIK VAN KINDERS Hou dig toe.</p> <p>REGISTRASIENOMMER 33/16.4/0285</p> <p>NAAM EN BESIGHEIDSADRES VAN DIE HOUER VAN DIE REGISTRASIESERTIFIKAAT iNova Pharmaceuticals (Pty) Ltd. Rileyweg 15e, Bedfordview 2007</p> <p>DATUM VAN PUBLIKASIE VAN DIE VOUBILJET 26 April 2002</p> <p>inova pharmaceuticals LT015503 LT049.08.17</p>