


<p>Skeduleringstatus [S4]</p> <p>Eiendomsnaam en doseervorm DISIPAL Tablette</p> <p>Samestelling Elke tablet bevat orfenadriënhidrochloried 50 mg. Ander bestanddele sluit in meliëstysel, Opadry® geel, polivinilpirolidoon, gesuiwerde talk, steariensuur.</p>	<p>Bevat suiker (laktosemonohidraat 20 mg en versiersuiker 20 mg).</p> <p>Farmakologiese klassifikasie A.5.4.1 (Preparate teen Parkinsonisme)</p> <p>Farmakologiese werking DISIPAL is 'n anticholinergiese middel wat op die sentrale senuweestelsel werk. Disipal verlig die simptome van Parkinsonisme en soortgelyke kondisies deur middel van die werking op die ventrokoudale streek van die diënselalon. DISIPAL het ook 'n ligte</p>	<p>euforiese effek.</p> <p>Indikasies Parkinsonisme; pre-Parkinsonisme; seniele en pre-seniele depressie; neuroleptiese sindroom.</p> <p>Kontraïndikasies Gloukoom; prostaathipertrufie.</p> <p>Die veiligheid gedurende swangerskap is nie vasgestel nie.</p> <p>Waarskuwings en spesiale voorsorgmaatreëls Bevat laktosemonohidraat. Pasiënte met die oorerflike kondisie van</p>	<p>galaktose intoleransie byvoorbeeld galaktosemie, Laplander laktase tekort, glukose-galaktose wanabsorpsie of fruktose intoleransie moet nie DISIPAL gebruik nie.</p> <p>Bevat laktosemonohidraat en versiersuiker wat die glukemiese beheer van pasiënte met diabetes mellitus kan beïnvloed.</p> <p>Dosis en gebruiksaanwysings</p> <p>1. Parkinsonisme; neuroleptiese sindroom: tussen een tablet drie maal per dag en twee tablette vier maal per dag.</p>	<p>2. Pre-Parkinsonisme; seniele- en pre-seniele depressie: een tablet twee maal per dag.</p> <p>3. Neuroleptiese sindroom: tot drie tablette drie maal per dag.</p> <p>Neuwe-effekte DISIPAL kan 'n ligte perifere anticholinergiese effek op sommige pasiënte hê, wat aanleiding mag gee tot droogheid van die mond en benewelde visie.</p> <p>Bekende simptome van oordosering en besonderhede van die</p>	<p>behandeling daarvan Atropienagtige effekte en euforie.</p> <p>Simptomatiese behandeling is aangedui.</p> <p>Identifikasie Geel, film-bedekte tablet.</p> <p>Aanbieding Amber glasbottels wat 28, 65, 84 of 100 tablette bevat. Pasiënt gereed pakke van 28, 56 of 84 tablette.</p>	<p>Bergingsaanwysings Bewaar teen of benede 30 °C.</p> <p>HOU BUITE BEREIK VAN KINDERS.</p> <p>Registrasiënommer C764 (Wet 101/1965).</p> <p>Naam en besigheidsadres van applikant iNova Pharmaceuticals (Pty) Ltd, Rileyweg 15E, Bedfordview</p> <p>Datum van publikasie van hierdie voulijet 24 Julie 1979</p>	<p>10 MM BLANK FOR GLUE AREA</p>	<table border="1"> <tr> <td data-bbox="3251 217 3419 392"> <p>A.5.4.1 Anti-Parkinsonism preparations. Each DISIPAL® tablet contains: Orphenadrine hydrochloride 50 mg, Lactose monohydrate 20 mg, Sugar icing 20 mg.</p> <p>Dosage and directions for use: See enclosed leaflet. Store at or below 30 °C. Keep out of reach of children.</p> </td> <td data-bbox="3419 217 3587 392"> <p>A.5.4.1 Anti-Parkinsonisme middels. Elke DISIPAL® tablet bevat: Orfenadriënhidrochloried 50 mg, Laktose monohidraat 20 mg, Versiersuiker 20 mg.</p> <p>Dosis en gebruiksaanwysings: Sien ingeslote voulijet. Bewaar teen of benede 30 °C. Hou buite bereik van kinders.</p> </td> <td data-bbox="3587 217 3783 392"> <p>A: C764 (Act 101/1965) 100 TABLETS/TABLETTE</p> <p>[S4] Disipal</p> <p>Nova Pharmaceuticals (Pty) Ltd Tel: +27 11 087 0000 www.inovapharma.co.za</p> </td> </tr> <tr> <td data-bbox="3251 392 3419 394"> <p>LA050.01.18 LA0112</p> </td> <td data-bbox="3419 392 3587 394"> <p>Schedulering Status: NS1</p> </td> <td data-bbox="3587 392 3783 394"> <p>Reg. No: 1454.1.0278 Co. Reg. No. 1952/001640/07</p> </td> </tr> </table>	<p>A.5.4.1 Anti-Parkinsonism preparations. Each DISIPAL® tablet contains: Orphenadrine hydrochloride 50 mg, Lactose monohydrate 20 mg, Sugar icing 20 mg.</p> <p>Dosage and directions for use: See enclosed leaflet. Store at or below 30 °C. Keep out of reach of children.</p>	<p>A.5.4.1 Anti-Parkinsonisme middels. Elke DISIPAL® tablet bevat: Orfenadriënhidrochloried 50 mg, Laktose monohidraat 20 mg, Versiersuiker 20 mg.</p> <p>Dosis en gebruiksaanwysings: Sien ingeslote voulijet. Bewaar teen of benede 30 °C. Hou buite bereik van kinders.</p>	<p>A: C764 (Act 101/1965) 100 TABLETS/TABLETTE</p> <p>[S4] Disipal</p> <p>Nova Pharmaceuticals (Pty) Ltd Tel: +27 11 087 0000 www.inovapharma.co.za</p>	<p>LA050.01.18 LA0112</p>	<p>Schedulering Status: NS1</p>	<p>Reg. No: 1454.1.0278 Co. Reg. No. 1952/001640/07</p>
<p>A.5.4.1 Anti-Parkinsonism preparations. Each DISIPAL® tablet contains: Orphenadrine hydrochloride 50 mg, Lactose monohydrate 20 mg, Sugar icing 20 mg.</p> <p>Dosage and directions for use: See enclosed leaflet. Store at or below 30 °C. Keep out of reach of children.</p>	<p>A.5.4.1 Anti-Parkinsonisme middels. Elke DISIPAL® tablet bevat: Orfenadriënhidrochloried 50 mg, Laktose monohidraat 20 mg, Versiersuiker 20 mg.</p> <p>Dosis en gebruiksaanwysings: Sien ingeslote voulijet. Bewaar teen of benede 30 °C. Hou buite bereik van kinders.</p>	<p>A: C764 (Act 101/1965) 100 TABLETS/TABLETTE</p> <p>[S4] Disipal</p> <p>Nova Pharmaceuticals (Pty) Ltd Tel: +27 11 087 0000 www.inovapharma.co.za</p>												
<p>LA050.01.18 LA0112</p>	<p>Schedulering Status: NS1</p>	<p>Reg. No: 1454.1.0278 Co. Reg. No. 1952/001640/07</p>												

<p>12MM BLANK FOR GLUE AREA</p>	<p>Scheduling status [S4]</p> <p>Proprietary name and dosage form DISIPAL Tablets</p> <p>Composition Each tablet contains 50 mg orphenadrine hydrochloride. Other ingredients include maize starch, Opadry® yellow, polyvinylpyrrolidone,</p>	<p>purified talc, stearic acid. Contains sugar (lactose monohydrate 20mg and sugar icing 20mg).</p> <p>Pharmacological classification A.5.4.1 (Anti – Parkinsonism Preparation)</p> <p>Pharmacological action DISIPAL is a centrally acting anti-cholinergic. Its site of action is the ventro-caudal region of the diencephalon. This</p>	<p>action results in DISIPAL's ability to alleviate the symptoms of Parkinsonism and similar conditions. DISIPAL also has a mild euphoriant action.</p> <p>Indications Parkinsonism; pre-Parkinsonism; senile and pre-senile depression; neuroleptic syndrome.</p> <p>Contraindications Glaucoma; prostatic hypertrophy. Safety in pregnancy has not been established.</p>	<p>Warnings and special precautions Contains lactose monohydrate. Patients with the rare hereditary conditions of galactose intolerance e.g. galactosaemia, Lapp lactase deficiency, glucose-galactose malabsorption or fructose intolerance should not take DISIPAL.</p> <p>Contains lactose monohydrate and sugar icing which may have an effect on the glycaemic control of patients with</p>	<p>diabetes mellitus.</p> <p>Dosage and directions for use</p> <p>1. Parkinsonism; Neuroleptic Syndrome: between one tablet three times a day and two tablets four times a day.</p> <p>2. Pre-Parkinsonism; senile and pre-senile depression: one tablet twice a day.</p> <p>3. Neuroleptic Syndrome: up to three tablets three times a day.</p>	<p>Side effects DISIPAL may in some patients manifest a mild peripheral anti-cholinergic action resulting in dryness of the mouth and blurring of vision.</p> <p>Known symptoms of overdosage and particulars of its treatment Atropine-like effects and euphoria. Symptomatic treatment is indicated.</p>	<p>Identification Yellow, film coated tablet.</p> <p>Presentation Amber glass bottles containing 28, 65, 84 or 100 tablets. Patient ready packs of 28, 56 or 84 tablets.</p> <p>Storage instructions Store at or below 30 °C. KEEP OUT OF REACH OF CHILDREN.</p>	<p>Registration number C764 (Act 101/1965).</p> <p>Name and business address of applicant iNova Pharmaceuticals (Pty) Ltd 15E Riley Road, Bedfordview</p> <p>Date of publication of this package insert 24 July 1979 KR-0mgs/KTK/25/409/98 LA050.01.18 LT0112 xxxxxxxx</p> 
---	---	---	--	--	---	---	--	---