

## Professional Information

### SCHEDULING STATUS

**S1**

### 1 NAME OF MEDICINE

Andolex, benzydamine hydrochloride 22,5 mg/15 ml solution.

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 15 ml contains

benzydamine hydrochloride 22.5 mg

Preservative:

methyl hydroxybenzoate 0.1 % m/v

Contains alcohol 10 % v/v

SUGAR FREE.

### 3 PHARMACEUTICAL FORM

A pleasant tasting, clear, green solution.

### 4 CLINICAL PARTICULARS

#### 4.1 Therapeutic indication

Symptomatic relief of painful inflammatory conditions of the mouth and throat including:

Traumatic conditions: Pharyngitis following tonsillectomy or after the use of nasogastric tube

Inflammatory conditions: Pharyngitis, aphthous ulcers and oral ulceration due to radiation therapy

Dentistry: For use after dental operations.

#### 4.2 Posology and method of administration

Oral Rinse: Rinse or gargle with 15 ml (approximately one tablespoonful) every 1½ to 3 hours as required for pain relief. The solution should be expelled from the mouth after use.

Spray: 5 to 10 sprays directly onto the painful or inflamed area and swallow gently. Repeat every 1½ to 3 hours as necessary.

Andolex should generally be used undiluted, but if "stinging" occurs, the rinse may be diluted with water.

Avoid contact with the eyes.

Elderly: No special usage recommendations are made for elderly patients.

#### 4.3 Contraindications

Sensitivity to benzydamine hydrochloride or to any of the components of this Andolex (see section 6.1).

Andolex is not recommended in children under the age of 12 years.

#### 4.4 Special warnings and precautions for use

Do not swallow. If a stinging or burning sensation occurs, Andolex may be diluted with water. Avoid contact with the eyes.

Uninterrupted treatment should not exceed 7 days, except under medical supervision.

Andolex contains methyl hydroxybenzoate which may cause allergic reactions (possibly delayed).

Effects on ability to drive and use machines:

Andolex has no or negligible influence on the ability to drive and use machines.

#### 4.5 Interaction with other medicines and other forms of interaction

See section 4.3 and section 4.4.

#### 4.6 Fertility, pregnancy and lactation

The safety of Andolex in pregnancy and lactation has not been established.

#### 4.7 Effects on ability to drive and use machines

Andolex has no or negligible effect on the ability to drive and use machines.

#### 4.8 Undesirable effects

##### Immune system disorders

Less frequent: Hypersensitivity reactions including urticaria, rash, laryngospasm, photodermatitis.

Frequency unknown:

Serious allergic reaction (anaphylactic shock), signs of which may include difficulty breathing, chest pain or chest tightness, and/or feeling dizzy/faint, severe itching of the skin or raised lumps on the skin, swelling of the face, lips, tongue and/or throat, and which may be potentially life-threatening.

##### Nervous system disorders

Less frequent: Light-headedness.

##### Gastrointestinal disorders

Frequency unknown: Nausea and vomiting

##### General disorders and administrative site conditions

Less frequent: Oral tissue numbness and stinging or burning sensation, disturbance of taste.

#### 4.9 Overdose

See section 4.8.

There is no specific antidote for benzydamine and should excessive quantities be ingested, the treatment should be symptomatic and supportive.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

A.16 Ear, nose and throat preparations

Benzydamine exerts a local anti-inflammatory and analgesic action by stabilising the cell membrane and inhibiting prostaglandin synthesis.

### **5.2 Pharmacokinetic properties**

When administered as a local application benzydamine has a low systemic absorption which reduces the potential of systemic side effects. Metabolism is mainly through oxidation, dealkylation and conjugation.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

- Glycerol
- saccharin sodium
- mouthwash flavour
- polysorbate
- quinoline yellow
- patent blue
- purified water

### **6.2 Incompatibilities**

None known.

### **6.3 Shelf life**

24 months.

### **6.4 Special precautions for storage**

Store at or below 25 °C. Protect from light. Do not leave the uncartoned bottle in direct sunlight.

**KEEP OUT OF REACH OF CHILDREN**

### **6.5 Nature and contents of container**

Oral Rinse: Bottles containing 200 ml and 2,5 L

Spray: Bottles containing 30 ml

### **6.6 Special precautions for disposal**

No special requirements.

## **7 HOLDER OF CERTIFICATE OF REGISTRATION**

iNova Pharmaceuticals (Pty) Ltd

15e Riley Road

Bedfordview

South Africa

## **8 REGISTRATION NUMBER**

Z/16/40

## **9 DATE OF FIRST AUTHORISATION**

Date of registration: May 1994

## **10 DATE OF REVISION OF THE TEXT**

15 January 2021.